SUMNER COUNTY SCHOOLS ALLERGY/ANAPHYLAXIS EMERGENCY ACTION PLAN & MEDICATION ORDER

Student:				DOB:			
School:							
History of Asthma (circle) YES or NO *if yes, the s		NO	*if yes, the student is at higher risk for severe reaction				
Extremely reactive to the fo THEREFORE:	llowing	allerç	gens: _				
	•			tely if the allergen was LIKELY eaten, for ANY symptoms.			
☐ If checked, give epinephrine immediately if the allergen was DEFINITELY eaten, even if NO symptoms are apparent.							

IS THIS STUDENT COMPETENT TO CARRY & SELF-ADMINISTER EMERGENCY MEDICATION (circle) YES

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

FOR ANY OF THE FOLLOWING:

SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hivesover body, widespread redness



Repetitive vomiting, severe diarrhea

①



Feeling something bad is about to happen. anxiety, confusion







OR A COMBINATION of symptoms

from different body areas.

1. INJECT EPINEPHRINE IMMEDIATELY.

- 2. Call 911. Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
 - Antihistamine
 - Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve.

MILD SYMPTOMS









NO

NOSE

Itchv or runny nose, sneezing

Itchv mouth

A few hives. mild itch

Mild nausea or

discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA FOLLOW THE DIRECTIONS BELOW:

- 1. Antihistamines may be given, if ordered by a healthcare provider.
- 2. Stay with the person; alert emergency contacts.
- 3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brandor Generic:						
Epinephrine Dose: ☐ 0.01 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM						
Antihistamine Brand or Generic:						
Antihistamine Dose:						
Other (e.g., inhaler-bronchodilator if wheezing):						

SUMNER COUNTY SCHOOLS ALLERGY/ANAPHYLAXIS EMERGENCY ACTION PLAN & MEDICATION ORDER

PARENT/GUARDIAN AKNOWLEDGMENTS & AUTHORIZATIONS:

• I want this plan implemented for my child.

Parent/Guardian Signature:

- I authorize the school nurse, or nurse program representative, to contact and receive additional information from the prescribing physician, regarding the student's health plan, as needed. I understand this information will only be shared with staff on a "need to know" basis.
- **If my student self-administers his/her epinephrine** I understand it is the responsibility of the parent/guardian to provide backup epinephrine, in the event the student loses or forgets their medication.
- If my child self-administers his/her epinephrine I agree to release the school district and school personnel from all claims of liability if my child suffers any adverse reactions from self-administration of epinephrine.
- If my child self-administers his/her epinephrine I understand it is my responsibility to review the following with my child:
 - Epinephrine must be with them at all times and never left unattended;
 - Ensure they know when (signs & symptoms) & how to use prescribed epinephrine;
 - He/she understands they must notify an adult in charge immediately, if epinephrine is used;
 - He/she will only use medication as prescribed and will never share with other students.

Emergency Phone:						
<u> </u>						
PHYSICIAN/HEALTHCARE PROVIDER SIGNATURE:						
Print:	Signature:	Date:				
LOCATION OF BACKUP MEDICATION (if none, indicate reason):						

If epinephrine given provide EMS with time and injector, if requested